AKADÉMIAI KIADÓ

Acta Veterinaria
Hungarica

68 (2020) 4, 421

DOI:

10.1556/004.2020.10000

© 2021 The Author(s)

ERRATUM

Erratum: Heart rate, heart rate variability, faecal glucocorticoid metabolites and avoidance response of dairy cows before and after changeover to an automatic milking system

VIKTOR JURKOVICH^{1*}, FRUZZSINA LUCA KÉZÉR^{2,3},
FERENC RUFF⁴, MIKOLT BAKONY¹, MARGIT KULCSÁR⁵ and
LEVENTE KOVÁCS^{2,3}

¹ Department of Animal Hygiene, Herd Health and Veterinary Ethology, University of Veterinary Medicine, István utca 2, H-1078 Budapest, Hungary

² MTA-SZIE Large Animal Clinical Research Group, Üllő Dóra-major, Hungary

³ Institute of Animal Husbandry, Faculty of Agricultural and Environmental Science, Szent István University, Gödöllő, Hungary

⁴ Department of Methodology, Hungarian Central Statistical Office, Budapest, Hungary

⁵ Department and Clinic of Reproduction, University of Veterinary Medicine, Budapest, Hungary

ERRATUM: Acta Veterinaria Hungarica 65 (2), (2017) pp. 301–313.

DOI: 10.1556/004.2017.029

In the article, Table 2 of the original paper should be replaced with the following Table 2:

Table 2. Areas under the curves (AUC) for heart rate and heart rate variability parameters calculated for premilking and milking periods¹

Item ²	Indicative of sympathetic/vagal tone	PREMILK ⁵			MILK ⁶		
		PARL ³	AMS ⁴	P value	PARL	AMS	P value
AUC _{HR} (beats)	Vagal tone	48.9 ± 12.0	25.9 ± 7.6	0.022	39.5 ± 7.8	34.5 ± 9.3	1.000
AUC _{HF} (n.u. × min)		–23.1 ± 13.6	–5.3 ± 7.3	0.025	–29.2 ± 13.4	–0.8 ± 4.9	0.015
AUC _{LF/HF}	Symp./parasymp. balance	12.6 ± 0.4	9.2 ± 0.1	0.033	20.7 ± 1.9	13.7 ± 1.4	0.026
AUC _{SD1} (ms × min)	Vagal tone	–9.4 ± 5.6	–2.2 ± 1.8	0.036	–10.5 ± 4.7	–4.0 ± 2.1	0.042
AUC _{SD2/SD1}	Symp. tone	14.8 ± 3.9	0.6 ± 2.49	<0.001	9.4 ± 3.8	–2.9 ± 2.6	0.033

¹ Descriptive statistics are based on means ± SEM of non-transformed data; ²HR = heart rate; HF = the high-frequency component [where n.u. = normalised units]; SD1 = standard deviation of instantaneous R–R variability measured from axis 1 in the Poincaré plot; SD2/SD1 = the ratio between SD2 (standard deviation of long-term continuous R–R variability measured from axis 2 in the Poincaré plot) and SD1; ³PARL = the period of conventional milking; ⁴AMS = the period of automatic milking system; ⁵PREMILK = the period of waiting in the holding pen before entering the parlot/milking unit; ⁶MILK = the period of milking (udder preparation, milking, waiting for exit).

*Corresponding author.

E-mail: jurkovich.viktor@univet.hu

Open Access. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium for non-commercial purposes, provided the original author and source are credited, a link to the CC License is provided, and changes – if any – are indicated.